



King's Research Portal

DOI:

[10.1016/j.jad.2018.02.006](https://doi.org/10.1016/j.jad.2018.02.006)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Lisiecka Ford, D. M., Tozer, D., Morris, R. G., Lawrence, A. J., Barrick, T. R., & Markus, H. S. (2018). Involvement of the reward network is associated with apathy in cerebral small vessel disease. *Journal of Affective Disorders*. <https://doi.org/10.1016/j.jad.2018.02.006>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal


Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Author's Accepted Manuscript

Involvement of the reward network is associated with apathy in cerebral small vessel disease

Danuta M. Lisiecka Ford, Daniel Tozer, Robin G. Morris, Andrew J. Lawrence, Thomas R. Barrick, Hugh S. Markus



PII: S0165-0327(17)32394-7
DOI: <https://doi.org/10.1016/j.jad.2018.02.006>
Reference: JAD9552

To appear in: *Journal of Affective Disorders*

Received date: 21 November 2017
Revised date: 25 January 2018
Accepted date: 12 February 2018

Cite this article as: Danuta M. Lisiecka Ford, Daniel Tozer, Robin G. Morris, Andrew J. Lawrence, Thomas R. Barrick and Hugh S. Markus, Involvement of the reward network is associated with apathy in cerebral small vessel disease, *Journal of Affective Disorders*, <https://doi.org/10.1016/j.jad.2018.02.006>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Involvement of the reward network is associated with apathy in cerebral small vessel disease

Danuta M. Lisiecka Ford¹, Daniel Tozer¹, Robin G. Morris², Andrew J. Lawrence², Thomas R. Barrick³, Hugh S. Markus¹

¹Stroke Research Group, University of Cambridge, Department of Clinical Neurosciences, Cambridge, UK

²King's College London, Institute of Psychiatry, Psychology and Neuroscience, London, UK

³St. Georges, University of London, Neurosciences Research Centre, London, UK

*Correspondence to: Danuta Lisiecka Ford

University of Cambridge, R3, Box 183, Addenbrooke's Biomedical Campus, Cambridge, CB2 0QQ, UK. Phone number: +44 1223216619; Fax number: +44 1223257288.

dml45@cam.ac.uk

Abbreviation; SVD, small vessel disease; MRI, magnetic resonance imaging; WMH, white matter hyperintensities; DTI, diffusion tensor imaging; VTA, ventral tagmental area; SCANS, St George's Cognition and Neuroimaging in Stroke; GDS, Geriatric Depression Scale; SPGR, spoiled gradient recalled echo; FSL, FMRIB Software Library; AAL, Automated Anatomical Labelling

Abstract

Introduction—

Apathy is a common yet under-recognised feature of cerebral small vessel disease (SVD), but its underlying neurobiological basis is not yet understood. We hypothesized that damage to the reward network is associated with an increase of apathy in patients with SVD.

Methods—

In 114 participants with symptomatic SVD, defined as a magnetic resonance imaging confirmed lacunar stroke and confluent white matter hyperintensities, we used diffusion tensor imaging tractography to derive structural brain networks and graph theory to determine network efficiency. We determined which parts of the network correlated with apathy symptoms. We tested whether apathy was selectively associated with involvement of the reward network, compared with two “control networks” (visual and motor).

Results–

Apathy symptoms negatively correlated with connectivity in network clusters encompassing numerous areas of the brain. Network efficiencies within the reward network correlated negatively with apathy scores; ($r=-0.344$, $p<0.001$), and remained significantly correlated after co-varying for the two control networks. Of the three networks tested, only variability in the reward network independently explained variance in apathetic symptoms, whereas this was not observed for the motor or visual networks.

Limitations–

The analysis refers only to cerebrum and not cerebellum. The apathy measure is derivative of depression measure.

Discussion–

Our results suggest that reduced neural efficiency, particularly in the reward network, is associated with increased apathy in patients with SVD. Treatments which improve connectivity in this network may improve apathy in SVD, which in turn may improve psychiatric outcome after stroke.

Keywords: apathy, reward, neural network, graph theory, small vessel disease, lacunar stroke

Introduction

Cerebral small vessel disease (SVD) affects the small arteries of the brain supplying the white matter and deep grey matter nuclei of the brain. It results in pathological changes in the brain tissue and characteristic radiological features best detected using magnetic resonance imaging (MRI), including lacunar infarcts, T2-white matter hyperintensities (WMH), cerebral microbleeds, and more diffuse white matter damage measured using diffusion tensor imaging (DTI). SVD results in a characteristic cognitive profile with early stage specific impairment in executive function and information processing speed (Lawrence et al., 2013). The psychological changes in SVD, it has been suggested, result from a disconnection syndrome with white matter tracts underlying complex subcortical-cortical circuits disrupted by the different neuropathologies, this is supported by recent findings from MRI structural network analysis (Lawrence et al., 2014).

In addition to cognitive impairment, apathy is a common affective feature of SVD (Brodaty et al., 2005). Apathy comprises lack of motivation, behavioural and cognitive retardation and reduced emotional reactivity and it results in diminished goal-oriented behaviour (Marin, 1996). It is a prominent affective symptom of many neurological diseases. The

neurobiological mechanisms underlying apathy in SVD are not yet fully understood. Some studies report the volume of WMH as a predictor of apathy in SVD, (Grool et al., 2012; Gupta et al., 2014; Nebes et al., 2001; Yao et al., 2015; Yao et al., 2009) but this has not been confirmed in further studies (Cosin et al., 2015; Grool et al., 2013). In a recent study apathy in SVD was correlated with decreased DTI fractional anisotropy in the cingulum, fornix and uncinate fasciculus, suggesting that white matter damage in these brain regions is more specifically causal (Hollocks et al., 2015). A recent review has identified more broadly diminished connectivity in frontal, subcortical and parietal areas as associated with apathy across many neurological and psychiatric conditions (Moretti and Signori, 2016). In SVD it has been hypothesized that reduction in white matter integrity affects the efficiency of reward networks which in turn results in apathy (Quattrocchi and Bestmann, 2014).

The reward network may play an important role in occurrence of apathy due to its role in the motivational process (Murayama et al., 2010). This is supported by behavioural evidence that reward insensitivity is positively correlated with apathy levels in stroke patients (Rochat et al., 2013). Moreover, dopamine plays an important role in the reward network, and methylphenidate, a stimulant whose action results in elevated levels of dopamine in the synaptic cleft, increases activation in the reward network in healthy individuals (Völlm et al., 2004). Methylphenidate has also been shown to reduce apathy after stroke (Martin et al., 1995; Spiegel et al., 2009).

The concept of a reward network has been extensively studied in animals, healthy individuals and individuals with depression, addiction and schizophrenia (Bracht et al., 2015; Cooper, 2002; Haber, 2008; Haber and Knutson, 2010; Lebreton et al., 2009; Mobbs et al., 2003; Nestler and Carlezon, 2006; Rolls, 2000; Sesack and Grace, 2010; Suk Lee et al., 2015). The consensus is that nucleus accumbens and its connections to ventral tegmental area (VTA),

amygdala complex, including associated temporal pole, and the prefrontal cortex (particularly medial and orbitofrontal cortex) are the core of the network associated with reward processing (Bracht et al., 2015; Cooper, 2002; Haber, 2008; Haber and Knutson, 2010; Lebreton et al., 2009; Mobbs et al., 2003; Nestler and Carlezon, 2006; Rolls, 2000; Sesack and Grace, 2010; Suk Lee et al., 2015). It is hypothesized that the network is supported by specific fibre bundles, including the cingulum, uncinated fasciculus and medial forebrain bundle and these fibres connecting the nucleus accumbens to the other core brain regions (Von Der Heide et al., 2013; Wu et al., 2016). The nucleus accumbens calculates the probability and magnitude of an expected reward outcome and decides whether to initialize action towards it (Cooper, 2002; Nestler and Carlezon, 2006; Sesack and Grace, 2010). The amygdala complex is associated with positive affect and reinforcement (Haber, 2008; Murray, 2007; Nestler and Carlezon, 2006) whereas the temporal pole relays information between the orbitofrontal cortex and amygdala while being involved in the processing of social rewards (Lebreton et al., 2009; Mobbs et al., 2003). The orbitofrontal cortex is involved in neural representation of primary and complex reinforcers and in controlling and correcting reward behaviour (Haber, 2008; Haber and Knutson, 2010; Lebreton et al., 2009; Nestler and Carlezon, 2006; Rolls, 2000). It is supported in its role by anterior cingulate cortex and medial frontal cortex which are associated with subjective rating of pleasure (Haber, 2008; Haber and Knutson, 2010; Nestler and Carlezon, 2006; Suk Lee et al., 2015).

In this study we use this overall framework to test whether network efficiency, derived from structural MR network analysis is negatively associated with apathy in SVD and whether the reward network plays a key role in mediating this association. To address these questions we apply DTI network analysis in patients with symptomatic SVD and use graph theory to provide a quantitative measure of network efficiency. This technique quantifies how efficient and well-connected the information exchange pathways are within a network. Thus we

determine which parts of the network were associated with apathy in SVD, and whether there was selective involvement of the reward network in the association with apathy, as compared with two other “experimental control” networks.

Materials and Methods

Authors indicate that they will not make their data available to other researchers.

Participants

One hundred and twenty one participants with symptomatic SVD were recruited between 2007 and 2010. They were from stroke services at three hospitals covering a geographically contiguous region of South London and recruited as part of the St George’s Cognition and Neuroimaging in Stroke (SCANS) study (Lawrence et al., 2013). SVD was defined as a clinical lacunar stroke syndrome (Bamford et al., 1991) with radiological evidence of a lacunar infarct in a brain region consistent anatomically with the clinical syndrome localized in white matter or deep grey matter nuclei and of the size smaller than 1.5 cm, in addition to confluent WMH of Fazekas grade 2 (early confluent) or higher (Fazekas et al., 1987). The inclusion criteria for SCANS was a Fazekas of 2 score or more as determined by the local clinician. All MRIs were then centrally reviewed blinded to clinical, MRI and cognitive details. The Fazekas scores used in the analysis were based on these blinded assessments. Both periventricular and deep WMH were separately rated and an overall score also generated which took into account both periventricular and deep WMH. Exclusion criteria were: any stroke mechanism other than SVD including intra/extra-cranial large artery stenosis >50%, cardio-embolic source, subcortical infarcts > 1.5cm in diameter as these are often embolic, or any cortical infarcts; history of major neurological or psychiatric condition excepting depression; non-fluent in English; not suitable for MRI; and unable to give informed consent.

SCANS is a longitudinal prospective study of MRI and cognition in SVD. For this analysis only the baseline measures were used. Out of all the 121 participants included in SCANS study at baseline, seven participants were excluded due to inadequate MRI data (acquisition difficulties or analysis pipeline failure) leaving 114 participants in the final analysis. The study was approved by a local research ethics committee of London–Wandsworth (07/Q0803/82) and all participants were able and provided written informed consent. The capacity of the participants to give informed consent was assessed via the review of their clinical status and by a researcher trained in taking informed consent. The study was registered: www.ukctg.nihr.ac.uk, study ID: 4577 and followed institutional guidelines. The study was conducted in accordance with the Helsinki Declaration as revised in 1989.

Apathy measure

The Geriatric Depression Scale (GDS) (Yesavage et al., 1983) was administered in full and apathy was measured using a subset of six items from this scale. These items were: ‘prefer to stay at home’, ‘avoid social gatherings’, ‘dropped activities and interests’, ‘find life very exciting’, ‘hard to start new projects’ and ‘full of energy’. Responses to these items are ‘yes’ or ‘no’, converted numerically and aggregated to provide an apathy score. The subset of items used have previously been shown to identify apathy, and allow distinction from depressive symptoms, in both healthy older adults (Adams et al., 2004) and SVD (Hollocks et al., 2015), with a meta-analysis concluding that apathy/withdrawal consistently appears as a single distinguishable factor in factorial analyses of GDS in various populations (Kim et al., 2013).

MRI acquisition

The MR acquisition has been described previously (Lawrence et al., 2013). Briefly, all the images were obtained in a 1.5T General Electric Signa HDxt MRI scanner (General Electric,

Milwaukee, WI, USA) with maximum gradient 33mTm^{-1} and a propriety head-coil. All sequences covered the entire brain with a total acquisition time of 45 minutes. Sequences relevant to the work described here were a T1-weighted sequence using a coronal spoiled gradient recalled echo (SPGR) with 176 1.1mm thick slices (field of view = $240 \times 240\text{mm}^2$, matrix = 256×192), TR = 11.5ms, TE = 5ms and flip angle of 18° . DTI images were also acquired with an axial single shot spin echo planar sequence comprising 55 slices without any gap (field of view = $240 \times 240\text{mm}^2$, matrix = 96×96), TR = 15600ms and TE = 93.4ms resulting in 2.5mm^2 isotropic voxels. First, four images without diffusion weighting ($b = 0\text{smm}^{-2}$) were followed by 25 diffusion-weighted ($b = 1000\text{smm}^{-2}$) images with gradient applied in non-collinear directions. Subsequently, another four $b = 0\text{smm}^{-2}$ images were obtained followed by diffusion weighted images with the negative of the 25 previously acquired directions.

Image pre-processing and network measures

The image analysis methods have been described in detail previously (Lawrence et al., 2014). Briefly, DTI images were eddy current corrected, and a diffusion tensor was fitted to the DTI signal using FMRIB Software Library (FSL) v4.1 (Jenkinson et al., 2012). Whole-brain deterministic diffusion tensor tractography was performed with streamlines terminating at angles of $\geq 40^\circ$ between principal eigenvectors in adjacent voxels or at $\text{FA} < 0.2$, which was the threshold relating to the ability of deterministic tensor tractography to give a reliable directional information. Network nodes were defined on the basis of the Automated Anatomical Labelling (AAL) atlas (Tzourio-Mazoyer et al., 2002). For each subject their T1-weighted image was co-registered to their averaged B0 DTI map using FSL FLIRT (Jenkinson et al., 2002; Jenkinson and Smith, 2001). A transformation between the T1-image and MNI space was calculated using Advanced Normalization Tools. These transformations

were subsequently applied to the AAL image to transform it into each subject's specific DTI space.

Graph theory describes the brain as nodes or vertices which represent grey matter regions and edges or lines connecting them which represent white matter connections. Such a method of representation helps to quantify mathematical properties of organization in complex neural networks (Bullmore and Sporns, 2009) by allowing to represent potential pathways of information-flow within the network. Thus 90 grey matter nodes were obtained for each subject. Network edges were defined as the presence of streamlines derived from tractography, directly connecting any pair of nodes. Both length (l) and number of streamlines (N) were taken into account when calculating the edge weighting (w_{ij}) so that the weight of each edge was proportional to the number of its streamlines and inversely proportional to the length of the streamlines. The equation was modified from Hagmann et al. (Hagmann et al., 2007) and was:

$$w_{ij} = \frac{1}{2} \sum_{m=1}^N \frac{1}{l_m}$$

Thus undirected weights of edges were obtained for the matrix of 90x90 nodes.

Statistical calculations

Network Based Statistics, which is a validated method to perform statistics on large networks, was used to perform the correlation analysis between apathy and connectivity weights of clusters of nodes (Zalesky et al., 2010). Only edges present in at least 30 subjects were taken into account. Mass univariate testing was performed using network based statistics and correlating weights of each edge with apathy levels; edges whose weights correlated with apathy scores at a level of $t = 2.5$ or higher (corresponding approximately to $p \leq 0.01$ in a two-tailed test) were selected as supra-threshold connections for further analysis. The network based statistics algorithm was then used to extract topological clusters of edges

and nodes deemed significantly correlated with apathy in SVD ($p < 0.05$). Thus, nodes and edges forming a network in which connectivity strength was correlated with apathy levels were identified.

We then investigated whether differences in reward network efficiency were related to variations in apathy. Efficiency of the network is defined as the average value of the shortest inverse path length between each pair of nodes participating in the network. The higher the efficiency of the network, the better-connected the information exchange pathways in the network are. If the efficiency of the network is high the information needs to travel via fewer stages when flowing from node to node. To test the specificity of any association we compared the associations to the degree of variance in apathy explained by two ‘experimental control’ networks, the motor and the visual networks. The three pre-defined sub-networks were delineated from the 90 AAL nodes obtained for each participant. Regions involved in the three networks were based on meta-analyses and review of papers of functional MRI and tractography studies in healthy individuals and individuals with depression (Bracht et al., 2015; Von Der Heide et al., 2013; Wu et al., 2016).

The networks were delineated as follows:

- a) Reward: left and right anterior cingulate cortex (frontal part of cingulum bundle), left and right putamen (nucleus accumbens incorporated), left and right inferior orbitofrontal cortex, left and right medial orbitofrontal cortex, left and right superior medial cortex, right and left amygdala, right and left superior temporal pole, right and left middle temporal pole (uncinate fasciculus)
- b) Motor: left and right precentral cortex, left and right supplementary motor area, and left and right putamen

- c) Visual: left and right superior occipital cortex, left and right middle occipital cortex, left and right inferior occipital cortex, left and right fusiform gyrus, and left and right parietal superior cortex

The efficiency of connectivity within each of these three networks was calculated using the weights of all possible edges between the nodes within the network, with the Brain Connectivity Toolbox (Rubinov and Sporns, 2010). The efficiency of each network was correlated with the apathy scores ($p < 0.05$ considered as significant). To test whether efficiency of any of the networks explained additional variance in apathy levels relative to the two other networks, we performed partial correlation between apathy and efficiency of each of the networks while controlling for efficiency in each of the other networks. The partial correlation was regarded as significant if the p-value was < 0.05 . Since the network efficiency is affected by possible disconnections, we repeated the analysis with the use of sum of weights instead of efficiency of the a-priori sub-networks. The same significance levels were applied in this analysis.

Controlling for age, vascular risk factors and SVD burden

Additionally, since age and vascular risk factors, that is hypertension (defined as systolic blood pressure $> 140\text{mmHg}$ or diastolic blood pressure $> 90\text{mmHg}$ or on hypertensive medication), hypercholesterolemia (defined as on statins therapies or a total cholesterol $> 5.2\text{mmol/l}$), diabetes mellitus (clinical diagnosis) and smoking (defined as current smoking), are important determinants of SVD we tested whether controlling for these risk factors changed the significance of the correlation between apathy and efficiency of the three neural networks. To achieve this goal we used partial correlation with the p-value < 0.05 deemed as significant.

We also tested whether apathy correlated with one of radiological markers of the severity of SVD. To do this we determined WMH volume, number of lacunas, number of cerebral microbleeds and number of perivascular spaces, using methods previously described (Benjamin et al., 2018; Benjamin et al., 2016; Lawrence et al., 2013). Briefly, WMH volume was calculated based on single-rater delineation of lesions on FLAIR images in the semi-automated DISPUNC program. Number of lacunas was determined on the basis of T1-weighted and FLAIR images, where a lacuna was defined as a CSF filled cavity within the white matter or subcortical regions, between 3–15 mm in diameter. Number of microbleeds was quantified on the basis of GRE images, where a microbleed was a well-defined focal area of low signal <10 mm in diameter. Number of perivascular spaces was calculated on the basis of T2 images where a perivascular space was defined as a smaller than 3mm punctuate or linear hyperintensity in basal ganglia or centrum semiovale. To do all this we performed correlations which were regarded significant if the p-value was <0.05.

Results

Demographic and clinical variables of the study population are shown in Table 1. The mean age was 70 years (SD=9.6 years), and 75/114 (65.8%) were male.

Figure 1 illustrates topological clusters in which connectivity strength correlated negatively with apathy. The clusters encompassed medial frontal lobes, basal ganglia, parietal lobes and temporal nodes. Table 2 presents the p-values of the correlation in each of the four observed clusters, the number of edges and nodes involved and the location of the nodes that participate in it. Some of the nodes observed in these clusters, such as the putamen, anterior cingulate cortex or inferior temporal pole, are considered to be part of the reward network, although the clusters also encompass other nodes which would not be classified as such, e.g. the precuneus or angular gyrus. In contrast to these negative correlations, there were no

positive correlations with apathy, suggesting that loss of connectivity is always associated with higher apathy.

The efficiencies of the three chosen sub-networks all correlated negatively with apathy scores; $r = -0.344$ for the reward network ($p < 0.001$), $r = -0.233$ for the motor network ($p = 0.013$) and $r = -0.210$ for the visual network ($p = 0.025$). The sums of weights of the three sub-networks were also all negatively correlated with the apathy scores: $r = -0.372$ for the reward network ($p < 0.001$), $r = -0.250$ for the motor network ($p = 0.007$) and $r = -0.210$ for the visual network ($p = 0.025$).

The partial correlation calculation showed that variability in the reward network efficiency explained additional variance in apathy when the correlation was controlled for efficiency in the other two networks. The reverse was not observed: that is, variability in efficiencies of the visual and motor network did not explain significant additional variance in apathy when the correlations were controlled for the effect of reward network. The partial correlation values and the corresponding p-values are presented in Table 3. The sums of weights within the sub-networks followed the same pattern, and their partial correlation values and corresponding p-values are also presented in Table 3.

The significance of the correlation between apathy and efficiency of the three neural networks did not change when age and vascular risk factors were controlled for. When controlled for these variables simultaneously apathy levels significantly correlated with efficiency of the reward network ($r = -0.355$, $p\text{-value} < 0.001$), motor network ($r = -0.234$, $p\text{-value} = 0.014$) and visual network ($r = -0.206$, $p\text{-value} = 0.031$).

Apathy did not correlate significantly with WMH volume ($r = 0.122$, $p\text{-value} = 0.198$), number of lacunas ($r = 0.169$, $p\text{-value} = 0.072$), number of microbleeds ($r = 0.139$, $p\text{-value} = 0.141$) or number of perivascular spaces ($r = 0.072$, $p\text{-value} = 0.445$).

In summary, this analysis shows that although the efficiencies of all three networks are mutually correlated, the efficiency of the reward network explains variability in apathy levels that is not explained by the other networks, risk factors or radiological markers of SVD severity.

Discussion

In our study we identified clusters of nodes whose connectivity is correlated with apathy in SVD. Among these nodes there were also some associated with reward processing. Subsequently, in our study we found that the efficiency of the reward network was negatively correlated with levels of apathy in individuals with SVD. Variability in reward network efficiency explained additional variance in apathy levels that was not explained by the efficiencies of two control networks, the motor and visual networks. This variability was also not explained by differences in age, vascular risk factors or severity of radiological markers of SVD. These results suggest white matter pathways underlying the reward networks are associated with increased levels of apathy in SVD.

The reward network has been previously associated with motivation and its disorders in healthy individuals and in patients (Barch et al., 2015; Bressan and Crippa, 2005; Suk Lee et al., 2015). Lesion studies in stroke suggest that damage to the structures pivotal for motivation and reward processing in humans, particularly the frontal lobes and basal ganglia, result in apathy (Grool et al., 2012; Grool et al., 2013; Gupta et al., 2014; Hama et al., 2011; Yang et al., 2013). Changes in white matter integrity have been previously associated with apathy in SVD (Hollocks et al., 2015). Our results provide further insight as to why apathy occurs to variable degrees in SVD. They not only emphasise the importance of white matter integrity, but also provide novel information on the specific neural localization and function of motivation and reward. Thus the results of this analysis suggest a united psychological and

neurobiological explanation of the phenomenon. They also emphasise a key role of the reward network in development or maintenance of apathy in SVD.

The findings may have implications for the treatment of apathy in SVD. Dopamine is an important neurotransmitter responsible for pharmacological connectivity within the human reward network. Increasing levels of dopamine could be a potential target for treatment of apathy in SVD. Initial pilot data towards such a treatment in patients suffering from stroke are promising, including medication such as methylphenidate or Ropinirole (Kohn et al., 2010; Martin et al., 1995; Spiegel et al., 2009).

Limitations

A potential limitation of the analysis is that all of the described network measures refer exclusively to the cerebrum, including the associated cerebral cortical and subcortical structures. The cerebellum was not included in the tractography analysis as there was a significant minority of participants whose field of view did not cover this region. Regions of the brain stem were also excluded because of the small size and a lack of T1 weighted images contrast relative to surrounding structures. Thus the nodes described here did not contain VTA which plays an important role in reward processing, nor parts of the cerebellum participating in motor control. In this paper we only test for the linear correlation between apathy and efficiency of the delineated networks. However, further exploration may be beneficial as to the linearity of the correlation.

Conclusion

In summary, this work implicates the importance of disruption of the reward network in determining whether patients with SVD suffer apathy. Treatments which improve connectivity in this network may improve apathy in SVD.

Author Statement**Contributors**

All the authors reviewed the manuscript and approved the final article. DMLF designed the analysis, performed and wrote the article. DT reviewed the analysis, contributed to it and reviewed the manuscript. RGM co-designed the original study and reviewed the manuscript. AJL collected data, contributed to the analysis and reviewed the manuscript. TRB co-designed the original study, reviewed the analysis and reviewed the manuscript. HSM co-designed the original study, reviewed the analysis and reviewed the manuscript.

Role of Funding Source

This work was supported by a Stroke Association Priority Programme Grant (PPA 2015/02). The SCANS study was supported by a grant from the Wellcome Trust (081589). HSM is supported by an NIHR Senior Investigator award. This work was supported by infrastructure provided by the Cambridge University Hospitals NIHR Comprehensive BRC.

Acknowledgements

The authors would like to thank Doctor Matthew Ford for editing of the manuscript.

Declaration of interest

None

Reference

Adams, K.B., et al., 2004. Confirmatory factor analysis of the geriatric depression scale. The Gerontologist 44, 818-826.

- Bamford, J., et al., 1991. Classification and natural history of clinically identifiable subtypes of cerebral infarction. *The Lancet* 337, 1521-1526.
- Barch, D.M., et al., 2015. Mechanisms underlying motivational deficits in psychopathology: similarities and differences in depression and schizophrenia, *Behavioral neuroscience of motivation*. Springer, pp. 411-449.
- Benjamin, P., et al., 2018. Lacunar infarcts, but not perivascular spaces, are predictors of cognitive decline in cerebral small vessel disease. *Stroke* In press.
- Benjamin, P., et al., 2016. Progression of MRI markers in cerebral small vessel disease: sample size considerations for clinical trials. *Journal of Cerebral Blood Flow & Metabolism* 36, 228-240.
- Bracht, T., et al., 2015. A review of white matter microstructure alterations of pathways of the reward circuit in depression. *Journal of affective disorders* 187, 45-53.
- Bressan, R.A., Crippa, J.A., 2005. The role of dopamine in reward and pleasure behaviour—review of data from preclinical research. *Acta Psychiatrica Scandinavica* 111, 14-21.
- Brodaty, H., et al., 2005. Frequency and clinical, neuropsychological and neuroimaging correlates of apathy following stroke—the Sydney Stroke Study. *Psychological medicine* 35, 1707-1716.
- Bullmore, E., Sporns, O., 2009. Complex brain networks: graph theoretical analysis of structural and functional systems. *Nature Reviews Neuroscience* 10, 186-198.
- Cooper, D.C., 2002. The significance of action potential bursting in the brain reward circuit. *Neurochemistry international* 41, 333-340.
- Cosin, C., et al., 2015. Circadian sleep/wake rhythm abnormalities as a risk factor of a poststroke apathy. *International Journal of Stroke* 10, 710-715.
- Fazekas, F., et al., 1987. MR signal abnormalities at 1.5 T in Alzheimer's dementia and normal aging. *American Journal of Neuroradiology* 8, 421-426.

- Grool, A., et al., 2012. Location and progression of cerebral small-vessel disease and atrophy, and depressive symptom profiles: The Second Manifestations of ARterial disease (SMART)-Medea study. *Psychological medicine* 42, 359-370.
- Grool, A.M., et al., 2013. Lacunar infarcts in deep white matter are associated with higher and more fluctuating depressive symptoms during three years follow-up. *Biological psychiatry* 73, 169-176.
- Gupta, M., et al., 2014. Behavioural and psychological symptoms in poststroke vascular cognitive impairment. *Behavioural Neurology* 2014, 1-5.
- Haber, S., 2008. Parallel and integrative processing through the Basal Ganglia reward circuit: lessons from addiction. *Biological psychiatry* 64, 173-174.
- Haber, S.N., Knutson, B., 2010. The reward circuit: linking primate anatomy and human imaging. *Neuropsychopharmacology* 35, 4.
- Hagmann, P., et al., 2007. Mapping human whole-brain structural networks with diffusion MRI. *PloS one* 2, e597.
- Hama, S., et al., 2011. Post-stroke depression and apathy: Interactions between functional recovery, lesion location, and emotional response. *Psychogeriatrics* 11, 68-76.
- Hollocks, M.J., et al., 2015. Differential relationships between apathy and depression with white matter microstructural changes and functional outcomes. *Brain* 138, 3803-3815.
- Jenkinson, M., et al., 2002. Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage* 17, 825-841.
- Jenkinson, M., et al., 2012. Fsl. *Neuroimage* 62, 782-790.
- Jenkinson, M., Smith, S., 2001. A global optimisation method for robust affine registration of brain images. *Medical image analysis* 5, 143-156.
- Kim, G., et al., 2013. A meta-analysis of the factor structure of the Geriatric Depression Scale (GDS): the effects of language. *Int Psychogeriatr* 25, 71-81.

- Kohno, N., et al., 2010. Successful treatment of post-stroke apathy by the dopamine receptor agonist ropinirole. *Journal of Clinical Neuroscience* 17, 804-806.
- Lawrence, A.J., et al., 2014. Structural network efficiency is associated with cognitive impairment in small-vessel disease. *Neurology* 83, 304-311.
- Lawrence, A.J., et al., 2013. Mechanisms of cognitive impairment in cerebral small vessel disease: multimodal MRI results from the St George's cognition and neuroimaging in stroke (SCANS) study. *PloS one* 8, e61014.
- Lebreton, M., et al., 2009. The brain structural disposition to social interaction. *European Journal of Neuroscience* 29, 2247-2252.
- Marin, R., 1996. Apathy: Concept, Syndrome, Neural Mechanisms, and Treatment, *Seminars in clinical neuropsychiatry*, pp. 304-314.
- Martin, E.M., et al., 1995. Successful methyphenidate treatment of apathy after subcortical infarcts. *Neurosciences* 7, 502-504.
- Mobbs, D., et al., 2003. Humor modulates the mesolimbic reward centers. *Neuron* 40, 1041-1048.
- Moretti, R., Signori, R., 2016. Neural Correlates for Apathy: Frontal-Prefrontal and Parietal Cortical- Subcortical Circuits. *Front Aging Neurosci* 8, 289.
- Murayama, K., et al., 2010. Neural basis of the undermining effect of monetary reward on intrinsic motivation. *Proceedings of the National Academy of Sciences* 107, 20911-20916.
- Murray, E.A., 2007. The amygdala, reward and emotion. *Trends in cognitive sciences* 11, 489-497.
- Nebes, R.D., et al., 2001. Relationship of deep white matter hyperintensities and apolipoprotein E genotype to depressive symptoms in older adults without clinical depression. *American Journal of Psychiatry* 158, 878-884.

- Nestler, E.J., Carlezon, W.A., 2006. The mesolimbic dopamine reward circuit in depression. *Biological psychiatry* 59, 1151-1159.
- Quattrocchi, G., Bestmann, S., 2014. Journal Club: possible role of the basal ganglia in poor reward sensitivity and apathy after stroke. *Neurology* 82, e171-e173.
- Rochat, L., et al., 2013. Poor reward sensitivity and apathy after stroke Implication of basal ganglia. *Neurology* 81, 1674-1680.
- Rolls, E.T., 2000. The orbitofrontal cortex and reward. *Cerebral cortex* 10, 284-294.
- Rubinov, M., Sporns, O., 2010. Complex network measures of brain connectivity: uses and interpretations. *Neuroimage* 52, 1059-1069.
- Sesack, S.R., Grace, A.A., 2010. Cortico-basal ganglia reward network: microcircuitry. *Neuropsychopharmacology* 35, 27.
- Spiegel, D.R., et al., 2009. Apathy due to cerebrovascular accidents successfully treated with methylphenidate: a case series. *The Journal of neuropsychiatry and clinical neurosciences* 21, 216-219.
- Suk Lee, J., et al., 2015. Neural basis of anhedonia and amotivation in patients with schizophrenia: the role of reward system. *Current neuropharmacology* 13, 750-759.
- Tzourio-Mazoyer, N., et al., 2002. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* 15, 273-289.
- Völlm, B.A., et al., 2004. Methamphetamine activates reward circuitry in drug naive human subjects. *Neuropsychopharmacology* 29, 1715-1722.
- Von Der Heide, R.J., et al., 2013. Dissecting the uncinate fasciculus: disorders, controversies and a hypothesis. *Brain* 136, 1692-1707.

- Wu, Y., et al., 2016. Segmentation of the cingulum bundle in the human brain: a new perspective based on DSI tractography and fiber dissection study. *Frontiers in Neuroanatomy* 10, 84.
- Yang, S.-r., et al., 2013. Predictors of early post ischemic stroke apathy and depression: a cross-sectional study. *BMC psychiatry* 13, 1.
- Yao, H., et al., 2015. Leisure-Time Physical Inactivity Associated with Vascular Depression or Apathy in Community-Dwelling Elderly Subjects: The Sefuri Study. *Journal of Stroke and Cerebrovascular Diseases* 24, 2625-2631.
- Yao, H., et al., 2009. Hypertension and white matter lesions are independently associated with apathetic behavior in healthy elderly subjects: the Sefuri brain MRI study. *Hypertension Research* 32, 586-590.
- Yesavage, J.A., et al., 1983. Development and validation of a geriatric depression screening scale: a preliminary report. *Journal of psychiatric research* 17, 37-49.
- Zalesky, A., et al., 2010. Network-based statistic: identifying differences in brain networks. *Neuroimage* 53, 1197-1207.

Figure 1. Topological clusters showing significant negative correlation between apathy and connectivity strength on the level $p < 0.05$ (corrected for multiple comparisons) in the whole brain analysis. Different colours represent the four clusters whose neural connectivity correlates negatively with apathy. The nodes forming the clusters are described in Table 2 (cluster 1 = red; cluster 2 = yellow; cluster 3 = green; cluster 4 = blue).

Table 1. Demographic and clinical variables of participants with SVD in the final analysis

Number of participants	114
Mean age in years (SD)	70.0 (9.6)

Gender male (%)	75 (65.8%)
Hypertension (%)	101 (88.6%)
Systolic BP (SD)	147.16 (21.5)
Diastolic BP (SD)	81.06 (10.7)
Hypercholesterolemia (%)*	104 (91.2%)
Diabetes mellitus (%)	22 (19.3%)
Current or ex-smoker (%)	63 (52.6%)
BMI kg/m ² (SD)	26.9 (4.8)
Fazekas Scale (%)	Fazekas 3 – 31 (27.2%)
	Fazekas 2 – 66 (57.9%)
	Fazekas 1 – 17 (14.9%)
Apathy (SD)	2.9 (1.7)

*Hypercholesterolemia defined as a previous diagnosis or total cholesterol above 5.2mmol/l.

Table 2. Nodes participating in the topological clusters showing a significant correlation between the connectivity strength and apathy levels in the whole brain analysis ($p < 0.05$).

Topological cluster	Edges count	Nodes count	Nodes participating	P-value
Cluster 1	13	13	L middle orbitofrontal cortex L&R medial superior frontal L anterior cingulate cortex L pallidum R middle cingulate cortex L hippocampus R thalamus L&R superior parietal cortex L&R precuneus R inferior parietal cortex	P<0.001
Cluster 2	9	9	L superior temporal pole L rolandic Operculum L superior temporal cortex L inferior temporal cortex L middle temporal cortex L angular gyrus L lingual gyrus	P=0.002

Cluster 3	6	7	L middle occipital cortex L inferior occipital cortex R superior temporal cortex R inferior temporal cortex R middle temporal cortex R angular cortex L calcarine fissure R middle occipital cortex R inferior occipital cortex	P=0.009
Cluster 4	4	5	R middle orbitofrontal cortex R inferior orbitofrontal cortex R superior temporal pole R insula R putamen	P=0.039

Table 3. Partial correlation between efficiency of a-priori defined sub-networks and apathy levels with the contribution of the remaining networks controlled

Network	Control network	Apathy correlation		p-value (two-tailed)	
correlated		value when controlled		when controlled for other	
		for other networks		networks	
		Efficiency	Sum of weights	Efficiency	Sum of weights
Reward network	Visual network	-0.289*	-0.323*	P=0.002	P<0.001
	Motor network	-0.27*	-0.293*	P=0.004	P=0.002
Motor network	Reward network	-0.074	-0.075	P=0.439	P=0.432
	Visual network	-0.177	-0.203*	P=0.06	P=0.031
Visual network	Reward network	-0.08	-0.051	P=0.401	P=0.588
	Motor network	-0.145	-0.136	P=0.125	P=0.151

*Significant partial correlations (p<0.05)

Highlights

- Apathy is an important affective consequence of cerebral small vessel disease, which may influence quality of life and adherence to rehabilitation
- Our study of 121 individuals with cerebral small vessel disease shows that neurobiological mechanism of apathy is associated with disconnectivity of white matter
- The white matter disconnectivity in the network responsible for reward processing is of particular importance for apathy symptoms in small vessel disease
- Treatments that target connectivity within the reward network may be effective in treating apathy after stroke

